

1968 AMC 12

Time limit: 75 minutes

Typeset by: LIVE by Po-Shen Loh

<https://live.poshenloh.com/past-contests/amc12/1968>



Copyright: Mathematical Association of America. Reproduced with permission.

1. Let P units be the increase in the circumference of a circle resulting from an increase of π units in the diameter. Then P equals:

- A $\frac{1}{\pi}$
- B π
- C $\frac{\pi^2}{2}$
- D π^2
- E 2π

2. The real value of x such that 64^{x-1} divided by 4^{x-1} equals 256^{2x} is:

A $-\frac{2}{3}$

B $-\frac{1}{3}$

C 0

D $\frac{1}{4}$

E $\frac{3}{8}$

3. A straight line passing through the point $(0, 4)$ is perpendicular to the line $x - 3y - 7 = 0$. Its equation is:

A $y + 3x - 4 = 0$

B $y + 3x + 4 = 0$

C $y - 3x - 4 = 0$

D $3y + x - 12 = 0$

E $3y - x - 12 = 0$

4. Define an operation $*$ for positive real numbers by $a * b = \frac{ab}{a+b}$. Then $4 * (4 * 4)$ equals:

A $\frac{2}{3}$

B 1

C $\frac{4}{3}$

D 2

E $\frac{16}{3}$

5. If $f(n) = \frac{1}{3}n(n+1)(n+2)$, then $f(r) - f(r-1)$ equals:

A $r(r+1)$

B $(r+1)(r+2)$

C $\frac{1}{3}r(r+1)$

D $\frac{1}{3}(r+1)(r+2)$

E $\frac{1}{3}r(r+1)(2r+1)$

6. Let side AD of convex quadrilateral $ABCD$ be extended through D , and let side BC be extended through C , to meet in point E . Let S represent the degree-sum of angles CDE and DCE , and let S' represent the degree-sum of angles BAD and ABC . If $r = \frac{S}{S'}$, then:

A $r = 1$ sometimes, $r > 1$ sometimes

B $r = 1$ sometimes, $r < 1$ sometimes

C $0 < r < 1$

D $r > 1$

E $r = 1$

7. Let O be the intersection point of medians AP and CQ of triangle ABC . If OQ is 3 inches, then OP , in inches, is:

A 3

B $\frac{9}{2}$

C 6

D 9

E undetermined

8. A positive number is mistakenly divided by 6 instead of being multiplied by 6. Based on the correct answer, the error thus committed, to the nearest percent, is:

- A 100
- B 97
- C 83
- D 17
- E 3

9. The sum of the real values of x satisfying $|x + 2| = 2|x - 2|$ is:

- A $\frac{1}{3}$
- B $\frac{2}{3}$
- C 6
- D $6\frac{1}{3}$
- E $6\frac{2}{3}$

10. Assume that, for a certain school, it is true that

I: Some students are not honest.

II: All fraternity members are honest.

A necessary conclusion is:

- ☐ A Some students are fraternity members
- ☐ B Some fraternity members are not students
- ☐ C Some students are not fraternity members
- ☐ D No fraternity member is a student
- ☐ E No student is a fraternity member

11. If an arc of 60° on circle I has the same length as an arc of 45° on circle II, the ratio of the area of circle I to that of circle II is:

- ☐ A 16 : 9
- ☐ B 9 : 16
- ☐ C 4 : 3
- ☐ D 3 : 4
- ☐ E none of these

12. A circle passes through the vertices of a triangle with side-lengths $7\frac{1}{2}$, 10 , $12\frac{1}{2}$. The radius of the circle is:

- A $\frac{15}{4}$
- B 5
- C $\frac{25}{4}$
- D $\frac{35}{4}$
- E $\frac{15\sqrt{2}}{2}$

13. If m and n are the roots of $x^2 + mx + n = 0$, $m \neq 0$, $n \neq 0$, then the sum of the roots is:

- A $-\frac{1}{2}$
- B -1
- C $\frac{1}{2}$
- D 1
- E undetermined

14. If x and y are nonzero numbers such that $x = 1 + \frac{1}{y}$ and $y = 1 + \frac{1}{x}$, then y equals:

A $x - 1$

B $1 - x$

C $1 + x$

D $-x$

E x

15. Let P be the product of any three consecutive positive odd integers. The largest integer dividing all such P is:

A 15

B 6

C 5

D 3

E 1

16. If x is such that $\frac{1}{x} < 2$ and $\frac{1}{x} > -3$, then:

A $-\frac{1}{3} < x < \frac{1}{2}$

B $-\frac{1}{3} < x < 3$

C $x > \frac{1}{2}$

D $x > \frac{1}{2}$ or $-\frac{1}{3} < x < 0$

E $x > \frac{1}{2}$ or $x < -\frac{1}{3}$

17. Let $f(n) = \frac{x_1 + x_2 + \cdots + x_n}{n}$, where n is a positive integer. If $x_k = (-1)^k, k = 1, 2, \dots, n$, the set of possible values of $f(n)$ is:

A $\{0\}$

B $\left\{\frac{1}{n}\right\}$

C $\left\{0, -\frac{1}{n}\right\}$

D $\left\{0, \frac{1}{n}\right\}$

E $\left\{1, \frac{1}{n}\right\}$

18. Side AB of triangle ABC has length 8 inches. Line DEF is drawn parallel to AB so that D is on segment AC , and E is on segment BC . Line AE extended bisects angle FEC . If DE has length 5 inches, then the length of CE , in inches, is:

- A $\frac{51}{4}$
- B 13
- C $\frac{53}{4}$
- D $\frac{40}{3}$
- E $\frac{27}{2}$

19. Let n be the number of ways that 10 dollars can be changed into dimes and quarters, with at least one of each coin being used. Then n equals:

- A 40
- B 38
- C 21
- D 20
- E 19

20. The measures of the interior angles of a convex polygon of n sides are in arithmetic progression. If the common difference is 5° and the largest angle is 160° , then n equals:

- ☐ A 9
- ☐ B 10
- ☐ C 12
- ☐ D 16
- ☐ E 32

21. If all the operations in $S = 1! + 2! + 3! + \cdots + 99!$ are correctly performed, the units digit in the value of S is:

- ☐ A 9
- ☐ B 8
- ☐ C 5
- ☐ D 3
- ☐ E 0

22. A segment of length 1 is divided into four segments. Then there exists a simple quadrilateral with the four segments as sides if and only if each segment is:

- ☐ A equal to $\frac{1}{4}$
- ☐ B equal to or greater than $\frac{1}{8}$ and less than $\frac{1}{2}$
- ☐ C greater than $\frac{1}{8}$ and less than $\frac{1}{2}$
- ☐ D greater than $\frac{1}{8}$ and less than $\frac{1}{4}$
- ☐ E less than $\frac{1}{2}$

23. If all the logarithms are real numbers, the equality

$$\begin{aligned}\log(x + 3) + \log(x - 1) \\ = \log(x^2 - 2x - 3)\end{aligned}$$

is satisfied for:

- ☐ A all real values of x
- ☐ B no real values of x
- ☐ C all real values of x except $x = 0$
- ☐ D no real values of x except $x = 0$
- ☐ E all real values of x except $x = 1$

24. A painting $18'' \times 24''$ is to be placed into a wooden frame with the longer dimension vertical. The wood at the top and bottom is twice as wide as the wood on the sides. If the frame area equals that of the painting itself, the ratio of the smaller to the larger dimension of the framed painting is:

A $1 : 3$

B $1 : 2$

C $2 : 3$

D $3 : 4$

E $1 : 1$

25. Ace runs with constant speed and Flash runs x times as fast, $x > 1$. Flash gives Ace a head start of y yards, and, at a given signal, they start off in the same direction. Then the number of yards Flash must run to catch Ace is:

A xy

B $\frac{y}{x + y}$

C $\frac{xy}{x - 1}$

D $\frac{x + y}{x + 1}$

E $\frac{x + y}{x - 1}$

26. Let $S = 2 + 4 + 6 + \cdots + 2N$, where N is the smallest positive integer such that $S > 1,000,000$. Then the sum of the digits of N is:

A 27

B 12

C 6

D 2

E 1

27. Let

$$S_n = 1 - 2 + 3 - 4 + \cdots + (-1)^{n-1}n.$$

Here $n = 1, 2, \dots$. Then $S_{17} + S_{33} + S_{50}$ equals:

A 0

B 1

C 2

D -1

E -2

28. If the arithmetic mean of a and b is double their geometric mean, with $a > b > 0$, then a possible value for the ratio $\frac{a}{b}$, to the nearest integer, is:

A 5

B 8

C 11

D 14

E none of these

29. Given the three numbers $x, y = x^x, z = x^{(x^x)}$, with $0.9 < x < 1.0$. Arranged in order of increasing magnitude, they are:

A x, z, y

B x, y, z

C y, x, z


D y, z, x

E z, x, y

30. Convex polygons P_1 and P_2 are drawn in the same plane with n_1 and n_2 sides, respectively, $n_1 \leq n_2$. If P_1 and P_2 do not have any line segment in common, then the maximum number of intersections of P_1 and P_2 is:

- A $2n_1$
- B $2n_2$
- C n_1n_2
- D $n_1 + n_2$
- E none of these

31. In this diagram, not drawn to scale, figures I and III are equilateral triangular regions with respective areas of $32\sqrt{3}$ and $8\sqrt{3}$ square inches. Figure II is a square region with area 32 square inches. Let the length of segment AD be decreased by $12\frac{1}{2}\%$ of itself, while the lengths of AB and CD remain unchanged. The percent decrease in the area of the square is:



- A $12\frac{1}{2}$
- B 25
- C 50
- D 75
- E $87\frac{1}{2}$

32. A and B move uniformly along two straight paths intersecting at right angles in point O . When A is at O , B is 500 yards short of O . In 2 minutes they are equidistant from O , and in 8 minutes more they are again equidistant from O . Then the ratio of A 's speed to B 's speed is:

A 4 : 5

B 5 : 6

C 2 : 3

D 5 : 8

E 1 : 2

33. A number N has three digits when expressed in base 7. When N is expressed in base 9 the digits are reversed. Then the middle digit is:

A 0

B 1

C 3

D 4

E 5

34. With 400 members voting, the House of Representatives defeated a bill. A re-vote, with the same members voting, resulted in passage of the bill by twice the margin by which it was originally defeated. The number voting for the bill on the re-vote was $\frac{12}{11}$ of the number voting against it originally. How many more members voted for the bill the second time than voted for it the first time?

A 75


B 60

C 50

D 45

E 20

35. In this diagram the center of the circle is O , the radius is a inches, chord EF is parallel to chord CD , O, G, H, J are collinear, and G is the midpoint of CD . Let K (square inches) represent the area of trapezoid $CDFE$ and let R (square inches) represent the area of rectangle $ELMF$. Then, as CD and EF are translated upward so that OG increases toward the value a , while JH always equals HG , the ratio $K : R$ becomes arbitrarily close to:



- A 0
- B 1
- C $\sqrt{2}$
- D $\frac{1}{\sqrt{2}} + \frac{1}{2}$
- E $\frac{1}{\sqrt{2}} + 1$

Solutions: <https://live.poshenloh.com/past-contests/amc12/1968/solutions>

